



THE STUBBORN BELLY FAT PROTOCOL



Table of Contents

- 1 Why belly fat is the last to go
- 2 The 4-pillar system
- 3 Training Structure
- 4 The belly fat loss food list
- 5 Supplements that actually help
- 6 Simple 4-week execution plan
- 7 How to track progress
- 8 End note


Who this is for: Anyone who has already lost weight but still struggles with stubborn belly fat, lower belly pouch, or a soft midsection.

Important truth: You cannot spot-reduce fat. This protocol works by combining fat loss, hormone optimization, stress control, steps, strength training, core tightening, and correct food choices.

Why Belly Fat Is the Last to Go

Belly fat is influenced by stress hormones, insulin, sleep, activity, and muscle mass. It is usually the last area to lean out.

- High cortisol (stress, poor sleep)
- Low or sub-optimal testosterone (in men)
- Insulin resistance
- Low daily steps
- Weak deep core muscles (TVA)
- Alcohol and liquid calories




The 4-Pillar System

Pillar 1:
Daily Steps



(10k–15k)

Pillar 2:
Strength Training



(3–4 days/week)

Pillar 3:
Short HIIT



(2–3 days/week)

Pillar 4:
Core Training



(3–4 days/week)

Training Structure (Simple & Effective)

- **Daily Steps:** 10,000–15,000 per day
- **Strength:** Squat, hinge, push, pull, lunge, press (3–4x/week)
- **HIIT:** 15–20 min short intervals (2x/week)
- **Core:** Planks, leg raises, dead bug, crunches, vacuum breathing

The Belly-Fat-Loss Food List

Base Rule: Eat high-protein, high-fiber, low-processed foods that control insulin and cortisol.

A) Proteins (Include in every meal):

- Eggs, egg whites
- Chicken, fish
- Paneer, tofu
- Curd / Greek yogurt
- Dal, rajma, chole
- Whey or plant protein

B) Vegetables (Unlimited):

- Spinach, methi, broccoli, cauliflower
- Cabbage, bottle gourd, ridge gourd
- Cucumber, tomato, capsicum, carrot
- Onion, mushrooms

C) Carbs (Control portions, mostly around workouts):

- Rice (preferably post-workout)
- Roti / chapati
- Oats, millets
- Sweet potato, potato
- Fruits (1–2 per day max)

D) Fats (Essential but portion-controlled):

- Nuts & seeds
- Ghee, olive oil, mustard oil
- Peanut butter
- Egg yolk
- Avocado

E) Foods that help reduce bloating & water retention:

- Cucumber, bottle gourd, pumpkin
- Curd / buttermilk
- Coconut water
- Banana (small)
- Leafy greens

F) Avoid or strictly limit:

- Sugar, sweets, desserts
- Bakery, biscuits, fried foods
- Alcohol, beer, cocktails
- Packaged & ultra-processed foods
- Fruit juices and sweet drinks



Supplements That Actually Help (Not Fat Burners)

Supplements do not burn belly fat. They fix deficiencies, improve hormones, sleep, and recovery.

Vitamin D3:



2000–4000 IU daily

Omega-3:



2–3 g EPA+DHA

Zinc:



15–30 mg daily

Magnesium glycinate:



300–400 mg at night

Protein powder:



if protein intake is low



Avoid:

Fat burners, detox teas, L-carnitine, CLA, ACV pills.

Simple 4-Week Execution Plan

- **Daily:** 10k–15k steps
- **Strength:** 3–4 days/week
- **HIIT:** 2 days/week
- **Core:** 3–4 days/week
- Follow **food list** + **supplement** support daily
- **Sleep 7–8 hours**, control stress

How to Track Progress

Waist measurement
every 2 weeks



Front & side photos
every 2 weeks



Steps & workout consistency



Do not rely only on scale weight




Final Message

Belly fat is the last to go. Follow this protocol honestly for 8–12 weeks and your waist and body shape will change.

- No hacks.
- No shortcuts.
- **Only smart execution and consistency.**





GET UP TO 50% OFF

ON FITELO'S WEIGHT LOSS PLANS

DOWNLOAD THE APP NOW